

## TECHNICAL REPORT

# Measurement of Fecal Steroids in the Black Rhinoceros (*Diceros bicornis*) Using Group-Specific Enzyme Immunoassays for 20-Oxo-Pregnanes

Franz Schwarzenberger, Kristina Tomášová, Dana Holečková,  
Bernd Matern, and Erich Möstl

*Inst. f. Biochemie and L. Boltzmann-Inst. f. Vet. med. Endokrinologie, University of Veterinary Medicine, Vienna, Austria (F.S., E.M.); Zoo Dvůr Kralové, Czechia (K.T., D.H.); and Zoo Frankfurt, Germany (B.M.)*

The metabolism and excretion of progesterone in different animal species results in several fecal 5-reduced progesterone metabolites (pregnanes), which in recent studies were quantified using progesterone antibodies. To increase the accuracy of fecal 20-oxo-pregnane evaluations in the black rhinoceros, enzyme immunoassays (EIA) using antibodies against 5 $\alpha$ -pregnane-3 $\beta$ -ol-20-one 3HS:BSA (5 $\alpha$ -20-one EIA) and 5 $\beta$ -pregnane-3 $\alpha$ -ol-20-one 3HS:BSA (5 $\beta$ -20-one EIA) were developed. The assays showed high crossreactivities with pregnanes containing a 20-oxo group and are referred to as group-specific; results of these assays were compared with an EIA using an antibody against 6HS-progesterone (4-ene-20-one EIA). Fecal samples of both subspecies of the black rhinoceros (*Diceros bicornis michaeli*, n = 5, and *Diceros bicornis minor*, n = 1) during pregnancy were collected 1–3 times/week. HPLC separation showed three major immunoreactive fecal 20-oxo-pregnane peaks; their elution profiles and different crossreactivities in the three EIAs provided strong evidence that these peaks are 5 $\alpha$ -pregnane-3,20-dione, 5 $\alpha$ -pregnane-3 $\alpha$ -ol-20-one, and 5 $\alpha$ -pregnane-3 $\beta$ -ol-20-one. Pregnane values in the pregnant animals continuously increased between months 3–7 and were significantly ( $P < 0.01$ ) elevated above the levels of nonpregnant animals (0.2  $\mu\text{g/g}$ ) by week 11. During months 6–13 concentrations in the 5 $\alpha$ -20-one and in the 5 $\beta$ -20-one EIA (5–11  $\mu\text{g/g}$ ) were significantly ( $P < 0.01$ ) higher than in the 4-ene-20-one EIA (1.5–3  $\mu\text{g/g}$ ). In conclusion, the immunoreactive fecal 20-oxo-pregnane metabolites in the black rhinoceros are determined more accurately with antibodies against pregnane-20-one-C3 conjugates, as compared with a progesterone antibody. © 1996 Wiley-Liss, Inc.

Received for publication May 10, 1995; revision accepted July 27, 1995.

Address reprint requests to Dr. Franz Schwarzenberger, Institut f. Biochemie, Vet. Med. Univ., Linke Bahngasse 11, A-1030 Vienna, Austria.

**Key words:** noninvasive, reproductive monitoring, pregnancy diagnosis, rhinoceros conservation, *Perrisodactyla*

## INTRODUCTION

The two subspecies of African black rhinoceros, the eastern (*Diceros bicornis michaeli*) and the southern (*Diceros bicornis minor*), are critically endangered due to continued poaching. The African population was reduced to less than 2,000 animals by the end of 1992 [Klös and Frese, 1993]. To save the remaining black rhinoceros population the current conservation strategies consist of conservation breeding in zoos [Klös and Frese, 1993], strengthening of antipoaching measures, and translocation to sanctuaries in Africa [Kock et al., 1995].

Successful breeding management may benefit from reliable reproductive monitoring. Reproductive data of rhinoceroses in recent studies were gained from the analysis of steroids in blood [Kock et al., 1991; Berkeley, 1994]; saliva [Kuckelkorn and Dathe, 1990; Czekala et al., in press]; urine [Kassam and Lasley, 1981; Kasman et al., 1986; Wagner, 1986; Ramsay et al., 1987, 1994; Hodges and Green, 1989; Hindle et al., 1992]; and feces [Schwarzenberger et al., 1993a, 1994, 1995b; Berkeley, 1994]. Furthermore, the metabolism of radioactively labeled steroid hormones was studied in a white rhinoceros [Hindle and Hodges, 1990], and ultrasonographic imaging was used to investigate the reproductive tract of rhinoceroses [Adams et al., 1991; Schaffer et al., 1994].

The most appropriate techniques for reproductive monitoring in nondomestic animals are the analysis of noninvasively collected fecal and urine samples. Recent investigations of fecal immunoreactive progesterone metabolites in several animal species used progesterone antibodies [Desaulniers et al., 1989; Messier et al., 1990; Wasser et al., 1991, 1994; Graham et al., 1993; Kirkpatrick et al., 1993; Monfort et al., 1993; Möstl et al., 1993; Schwarzenberger et al., 1993a, 1995a; Berkeley, 1994; Brown et al., 1994; Czekala et al., 1994; Larter et al., 1994]. Resulting in useful biological data, most of these studies did not characterize the immunoreactive material by any method other than RIA. However, metabolism and fecal excretion of radioactively labeled progesterone in feline, primate, and farm animal species resulted in several progesterone metabolites [Shideler et al., 1993; Brown et al., 1994; Wasser et al., 1994; Palme, personal communication]. In addition, HPLC separation with subsequent immunoassay analysis displayed the presence of up to eight well defined immunoreactive fecal progesterone metabolites [Kirkpatrick et al., 1991; Schwarzenberger et al., 1991, 1992, 1993a,b, 1995a; Heistermann et al., 1993; Möstl et al., 1993; Shideler et al., 1993; Brown et al., 1994; Wasser et al., 1994]. These metabolites showed elution patterns similar to that of pregnanes, and "authentic" progesterone was either not detectable or detectable to a limited extent only.

Progesterone is metabolized to pregnanes before its fecal excretion. Some of these pregnanes (namely 5 $\alpha$ - and 5 $\beta$ -pregnane-3,20-dione) still possess two oxo groups in position C3 and C20 of the steroid molecule, and therefore these metabolites show crossreactivities with progesterone antibodies. However, depending on crossreactivities, specific progesterone antibodies quantify only a small proportion of the actual amount of fecal pregnane compounds. A possibility for determining these metabolites more accurately is the application of antibodies against pregnane-C3 conjugates. Since the functional group of the steroid in position C3 is masked, these

antibodies show considerable crossreactivities with several progesterone metabolites sharing a similar C20 configuration (a 20-oxo, a 20 $\alpha$ -OH, or a 20 $\beta$ -OH group) and their measurement is combined in one assay, which we refer to as group-specific.

In our recent study on the black rhinoceros we (1) determined the estrous cycle length as 25 days; (2) diagnosed pregnancy by means of increasing fecal immunoreactive progesterone metabolites after the first quarter of gestation; and (3) demonstrated the presence of immunoreactive 20-oxo- and 20 $\alpha$ -OH-pregnane compounds [Schwarzenberger et al., 1993a]. However, lacking specific enzyme immunoassays (EIA) for 20-oxo-pregnanes, we relied on the crossreactivities of an antibody against 6HS-progesterone, and assay specificity and accuracy were a serious concern. Therefore we raised antibodies against 5 $\alpha$ - and 5 $\beta$ -pregnanes containing a 20-oxo group and developed EIAs.

The aims of the present study were: (1) to develop specific assays for pregnanes containing a 20-oxo group (20-oxo-pregnanes); (2) to compare EIA measurements of fecal 20-oxo-pregnanes in three EIAs with different degrees of specificity; (3) to characterize the possible structure of the fecal immunoreactive 20-oxo-pregnanes; and (4) to determine fecal 20-oxo-pregnane values in both subspecies of the black rhinoceros.

## METHODS

### Animals and Sample Collection

Female black rhinoceroses of the eastern (*Diceros bicornis michaeli*) and the southern (*Diceros bicornis minor*) subspecies were studied. Animals of the eastern subspecies ( $n = 5$ ) were housed at Dvůr Kralové Zoo. At the times of parturitions reported, the animals were 10, 15, 16, and 23 years old and had (including deliveries reported in this study) 2, 3, 4, and 1 offspring, respectively. One 24-year-old animal had a late pregnancy abortion after 14 months of gestation; this was her fifth pregnancy and her previous pregnancies delivered viable calves. The rhino ( $n = 1$ ) of the southern subspecies was housed at Frankfurt Zoo; she was seven years old and the report in this study was her first pregnancy. Each female was allowed access to a male during the daytime and housed alone at night. Estrous behavior and mating were recorded by the keeper staff. Animals were fed hay and grass, supplemented with leaves, branches, fruits, and vegetables, and had free access to water.

Morning fecal samples were collected off the ground and stored at  $-20^{\circ}\text{C}$  until analysis. Samples were collected 2–3 times/week for the first 4 months of pregnancy and the month before parturition. Samples during midgestation were collected once a week or biweekly. Samples from nonpregnant animals were collected three times/week.

### Fecal Extraction for Analysis With EIAs

Fecal samples (0.5 g) were extracted with methanol as described previously [Schwarzenberger et al., 1993a], except 1.0 g powdered aluminum oxide was added before extraction. Briefly, 0.5 g feces was mixed with 1.0 g powdered aluminium oxide, extracted with 4.5 ml aqueous methanol (80%), and defatted with petroleum benzene. The methanol was diluted and analyzed with the EIAs. Aluminium oxide was added to remove visible background pigment [Graham et al., 1993] and in an

TABLE 1. Enzyme immunoassays (EIA)

EIA	4-ene-20-one <sup>a</sup>	5 $\alpha$ -20-one	5 $\beta$ -20-one
Immunogens	4-pregnene-6 $\alpha$ -ol- 20-one-6HS:BSA	5 $\alpha$ -pregnane-3 $\beta$ -ol- 20-one-3HS:BSA	5 $\beta$ -pregnane-3 $\alpha$ -ol- 20-one-3HS:BSA
Labels	5 $\alpha$ -pregnane-3 $\beta$ -ol- 20-one-3HS:DADOO-B <sup>b</sup>	4-pregnene-3,20-dione- 3CMO:DADOO-B <sup>b</sup>	5 $\alpha$ -pregnane-3 $\beta$ -ol- 20-one-3HS:DADOO-B <sup>b</sup>
Antibody titers	1:40 $\times$ 10 <sup>3</sup>	1:15 $\times$ 10 <sup>3</sup>	1:15 $\times$ 10 <sup>3</sup>
Label titers	1:250 $\times$ 10 <sup>3</sup>	1:600 $\times$ 10 <sup>3</sup>	1:600 $\times$ 10 <sup>3</sup>
Standard curve range	2–500 pg	2–500 pg	5–1,250 pg
Sensitivity (90% binding)	1.5 $\pm$ 0.12 pg	2.6 $\pm$ 0.33 pg	5.7 $\pm$ 0.37 pg
Intrassay coefficient of variation	10.4%	11.3%	9.6%
Interassay coefficient of variation	12.8%	16.2%	14.6%

<sup>a</sup>Assay described by Schwarzenberger et al. [1993].

<sup>b</sup>DADOO-B = biotin-DADOO = N-biotinyl-1,8-diamino-3,6-dioxaoctan.

initial study was found to increase differences between low and high 20-oxo-pregnane values in all three assays used.

### Analysis With EIAs

Fecal extracts were analyzed with two newly developed EIAs (Tables 1,2) and the data compared with that using the 4-ene-20-one EIA described in our previous study [Schwarzenberger et al., 1993a]. The antibodies for the three EIAs were raised in rabbits. Progesterone was used as standard, and serial dilutions of fecal extracts gave displacement curves parallel to that of the standard curve in each assay.

The three assays followed the protocol described by Schwarzenberger et al. [1991, 1993a], except that steroid-DADOO-biotin conjugates (DADOO-biotin = N-biotinyl-1,8-diamino-3,6-dioxaoctane) were used as labels [Palme and Möstl, 1993; Pryce et al., 1994]. Briefly, the three EIAs used a double-antibody technique and were performed on anti-rabbit-IgG-coated microtiter plates. After overnight incubation of standard or samples, steroid antibody, and biotinylated label, the plates were emptied, washed, and blotted dry, before a streptavidin horseradish peroxidase conjugate was added. After a 45 min incubation, plates were emptied, washed, and blotted dry. The substrate (tetramethylbenzidine) was added and incubated for 45 min at 4°C before the enzymatic reaction was stopped with 2 mol/l sulfuric acid.

The synthesis of the steroid-DADOO-biotin conjugates followed the mixed anhydride reaction [Kellie et al., 1975]. Steroid:HS or steroid:CMO derivatives (5–10 mg) were dissolved in 1 ml dimethylformamide, and cooled to –18°C under constant stirring. Then 0.01 ml methylmorpholin and 0.01 ml isobutylchloroformiate were added and incubated at –18°C for 30 min. Thereafter, 1.5 ml of a mixture of 0.1 mol/l sodium phosphate buffer (pH 8.2) and dimethylformamide (1/2 v/v; –18°C), containing 5 mg biotinyl-DADOO, was slowly added under constant readjustment of the pH, using 1 mol/l KOH. The mixture was incubated at room temperature for 3 hr, under stirring and light protection. Then 7.5 ml of water was added and the solution was passed through a primed C18 cartridge. The cartridge was washed with 5 ml water, and the steroid and steroid conjugates were eluted with 4 ml methanol. The methanol was evaporated (40°C) under nitrogen and the residue dis-

**TABLE 2. Crossreactivities (%) of steroids in the different enzyme immunoassays**

Steroid	4-ene-20-one	5 $\alpha$ -20-one	5 $\beta$ -20-one
4-pregnen-			
3,20 dione (progesterone)	100	100	100
3 $\beta$ -ol-20-one	26	14	68
3 $\alpha$ -ol-20-one	8	390	20
20 $\alpha$ -ol-3-one	<0.1	<0.1	<0.1
20 $\beta$ -ol-3-one	<0.1	<0.1	<0.1
5 $\alpha$ -pregnane-			
3 $\alpha$ -ol-20-one	5	89	8
3 $\beta$ -ol-20-one	18	56	102
3,20-dione	40	168	75
3 $\alpha$ ,20 $\alpha$ -diol	<0.1	<0.1	<0.1
3 $\alpha$ ,20 $\beta$ -diol	<0.1	<0.1	<0.1
5 $\beta$ -pregnane-			
3 $\alpha$ -ol-20-one	3	88	20
3 $\beta$ -ol-20-one	7	5	36
3,20-dione	71	26	151
3 $\alpha$ ,20 $\alpha$ -diol	<0.1	<0.1	<0.1
3 $\alpha$ ,20 $\beta$ -diol	<0.1	<0.1	<0.1
5-pregnen-			
3 $\beta$ -ol-20-one	29	17	11
Cortisol	<0.1	<0.1	<0.1
Estrone	<0.1	<0.1	<0.1

solved in 2 ml of 20% methanol. The solution was injected onto a preparative RP18 chromatography system (310 × 25 mm, flow rate 1 ml/min). Separation of the steroid-biotin-DADOO conjugate from the free steroids was achieved using a methanol gradient, linearly increasing from 20 to 100% between fractions 0–70. Fraction size was 5 ml, and the steroid-biotin-DADOO conjugate eluted between fractions 65–70.

### High Performance Liquid Chromatography (HPLC) of Fecal 20-oxo-pregnanes

The number, their relative proportions, and the polarity of the unconjugated fecal immunoreactive 20-oxo-pregnanes were determined by HPLC and subsequent analysis with the three respective EIAs [Schwarzenberger et al., 1991, 1993a]. Samples from both subspecies during different stages of pregnancy were used. Samples were mixed with <sup>3</sup>[H]-progesterone and <sup>3</sup>[H]-20 $\alpha$ -dihydroprogesterone, and extracted with methanol. The methanol was evaporated, the residue reconstituted in 1 ml of 20 mmol NaHCO<sub>3</sub>, and twice re-extracted with 5 ml of petroleum ether/diethylether (6/4 v/v). The ether was evaporated and the samples reconstituted in chloroform/n-hexane. Removal of polar substances was compatible with our straight phase HPLC system (Si 60, 10  $\mu$ m; 250 × 4 mm; flow rate 2 ml/min); however, it is possible that conjugated steroids were removed.

The HPLC separation followed the principles described previously [Schwarzenberger et al., 1991, 1993a], but the elution time between <sup>3</sup>[H]-progesterone and <sup>3</sup>[H]-20 $\alpha$ -dihydroprogesterone was increased by using chloroform/n-hexane (v/v 75/25) and a linear methanol gradient (0–6%). Retention times of the immunoreactive

fractions were compared with those of different crossreacting steroid standards ( $5\alpha/5\beta$ -pregnanes) and with  $^3\text{[H]}$ -progesterone and  $^3\text{[H]}$ - $20\alpha$ -dihydroprogesterone. Elution patterns of pregnane standards were determined by EIAs, and  $^3\text{[H]}$ -steroids were determined by liquid scintillation counting.

### Data Analysis

Data are presented as mean  $\pm$  SEM. A total of 216 samples from nonpregnant animals ( $n = 5$ ) during the estrous cycle was used for comparison with those of pregnant animals. The duration of pregnancy was calculated as the period between observed mating and parturition. Samples from the first 13 months and the last 2 months of pregnancy were standardized to the day of mating and to the day of parturition, respectively. Results were grouped in weekly intervals during gestational weeks 1–17 and 61–65, and in monthly intervals between these periods. Means calculated represent 5–18 samples per point. Statistical significance of the differences between pregnant and nonpregnant animals was tested by the Student's  $t$  test and the Mann–Whitney  $U$  test. Differences among the EIAs during pregnancy were analyzed by a paired  $t$  test and the Wilcoxon's signed-rank test.

## RESULTS

### Enzyme Immunoassays (EIA)

Characteristics of the EIAs are summarized in Tables 1 and 2. Antibodies for the  $5\alpha$ - $20$ -one and  $5\beta$ - $20$ -one EIA were raised against steroid-C3-hemisuccinate protein conjugates; thus, group-specific antibodies directed against the C20 position of the steroid were obtained. The assays quantify closely related 5-reduced progesterone metabolites (pregnanes) containing an oxo group on the C20 atom, and the results are expressed as 20-oxo-pregnanes. Crossreactivities with pregnanes were considerably higher in the  $5\alpha$ - $20$ -one and  $5\beta$ - $20$ -one EIA, as compared with the 4-ene- $20$ -one EIA. However, the former two EIAs did not completely differentiate between the  $5\alpha$ - and the  $5\beta$ -pregnane series. Either EIA could be used for fecal steroid monitoring, although HPLC separation provided strong evidence for the sole presence of  $5\alpha$ -pregnanes.

### HPLC Separation of Fecal 20-oxo-pregnanes

HPLC separation and subsequent analysis of the fractions with the three EIAs were performed on fecal samples ( $n = 5$ ) from both subspecies during different stages of pregnancy. A representative midpregnancy profile is shown in Figure 1. Results did not differ between the two subspecies; three major immunoreactive 20-oxo-pregnane peaks were detected in each sample. Their relative proportions remained unchanged throughout pregnancy. Immunoreactive material coeluting with  $^3\text{[H]}$ -progesterone was not detectable.

Comparison with corresponding reference steroids together with the different crossreactivities in the three respective EIAs provided good evidence that the fecal immunoreactive peaks are (1)  $5\alpha$ -pregnane-3,20-dione, (2)  $5\alpha$ -pregnane- $3\alpha$ -ol- $20$ -one, and (3)  $5\alpha$ -pregnane- $3\beta$ -ol- $20$ -one. Further evidence for the identity of peaks 2 and 3 was gained from the chromatography of fecal samples spiked with the authentic steroid standards, and after ether extraction and rechromatography of the fractions containing peaks 2 and 3. These separations resulted in chromatograms similar to that




Fig. 1. HPLC separation of immunoreactive 20-oxo-pregnanes in a fecal sample of a 7-months-pregnant black rhinoceros (*Diceros bicornis michaeli*). Elution patterns of (1)  $^3\text{H}$ -progesterone, (2)  $^3\text{H}$ -20 $\alpha$ -dihydroprogesterone were determined by liquid scintillation counting ( $\Delta$  cpm/fraction). Fractions were analyzed with EIAs for 20-oxo-pregnanes ( $\circ$  5 $\alpha$ -20-one EIA;  $\bullet$  5 $\beta$ -20-one EIA;  $\square$  4-ene-20-one EIA). Concentrations in ng/fraction were calculated for 1 g of feces without correction for methodological losses. Crossreactivities of immunoreactive peaks (a–c) in the three EIAs are given in the inset. Retention times of peaks a–c were comparable to: (a) 5 $\alpha$ -pregnane-3,20-dione, (b) 5 $\alpha$ -pregnane-3 $\alpha$ -ol-20-one, and (c) 5 $\alpha$ -pregnane-3 $\beta$ -ol-20-one.

obtained after fecal sample separation. The elution patterns of crossreacting 5 $\beta$ -pregnanes were different from the fecal immunoreactive metabolites; the steroid standards 5 $\beta$ -pregnane-3,20-dione, 5 $\beta$ -pregnane-3 $\alpha$ -ol-20-one, and 5 $\beta$ -pregnane-3 $\beta$ -ol-20-one eluted in fractions 10, 39–42, and 42–46, respectively (Fig. 1).

### Fecal 20-Oxo-Pregnane Profiles

Estrous cycle profiles were comparable to those described previously [Schwarzenberger et al., 1993a]. Distinction between follicular and luteal phases was possible in the three EIAs used, and mating corresponded with low fecal 20-oxo-pregnane concentrations. Therefore, mean values were calculated from samples collected during follicular and luteal phases, and used for comparison only with the results obtained during pregnancy. Values of the nonpregnant animals were  $117 \pm 8$ ,  $136 \pm 8$ , and  $140 \pm 9$  ng/g feces in the 4-ene-20-one, 5 $\alpha$ -20-one, and 5 $\beta$ -20-one EIA, respectively.

Gestation lengths for the animals of the *Diceros bicornis michaeli* subspecies were 442, 455, 467, and 469 days, respectively; one animal had a late pregnancy abortion on day 427 (Fig. 2). Values of this animal were used for mean  $\pm$  SEM calculation during the first 9 months of gestation only. Duration of pregnancy was 464 days for the animal of the *Diceros bicornis minor* subspecies.




Fig. 2. Concentrations of immunoreactive 20-oxo-pregnanes in fecal samples of a black rhinoceros (*Diceror bicornis michaeli*) during an estrous cycle, subsequent pregnancy, and after late pregnancy abortion. Samples were analyzed with EIAs for 20-oxo-pregnanes (○ 5 $\alpha$ -20-one EIA; ● 5 $\beta$ -20-one EIA). Note breaks of horizontal and vertical axes.

Pregnane concentrations during pregnancy were analyzed with the three EIAs and results of the *Diceror bicornis michaeli* (mean  $\pm$  SEM) and the *Diceror bicornis minor* subspecies are shown in Figures 3 and 4, respectively; endocrine profiles of the two subspecies were comparable. Mean 20-oxo-pregnane values increased to luteal phase values within 2 weeks after mating and remained in that range during the first 2 months of pregnancy. Pregnane concentrations continuously increased during months 3–7, and values were significantly ( $P < 0.01$ ) elevated above mean values of non-pregnant animals from week 11 onwards. The increase in the 4-ene-20-one EIA values was moderate as compared with the pronounced increase in the 5 $\alpha$ -20-one and 5 $\beta$ -20-one EIA. During months 6–13 mean values in the 5 $\alpha$ -20-one and the 5 $\beta$ -20-one EIA (5–11  $\mu\text{g/g}$ ) were significantly higher ( $P < 0.01$ ) than those in the 4-ene-20-one EIA (1.5–3  $\mu\text{g/g}$  feces). The 20-oxo-pregnane values decreased during the last 3 months of gestation and returned to basal levels within 2–3 days after parturition.

## DISCUSSION

The principal objective of this study was to compare measurements of fecal 20-oxo-pregnane values in EIAs with different degrees of specificity. Results of our recent study on fecal immunoreactive progesterone metabolites in the black rhinoceros (*Diceror bicornis*) demonstrated the presence of 20-oxo- and 20 $\alpha$ -OH-pregnanes; progesterone was not present, or present to a limited extent only [Schwarzenberger et al., 1993a]. However, lacking specific assays for 20-oxo-pregnanes, we relied on the crossreactivities of an antibody against 6HS-progesterone (the 4-ene-20-one EIA used in this study). Thus, assay specificity and accuracy of fecal 20-oxo-pregnane determinations were a reasonable concern. Therefore, two antibodies against 5 $\alpha$ - and 5 $\beta$ -pregnane-20-one 3HS:BSA derivatives (5 $\alpha$ -20-one EIA and 5 $\beta$ -






Fig. 3. Concentrations (mean  $\pm$  SEM;  $n = 5-18$  per point) of immunoreactive 20-oxo-pregnanes in fecal samples of black rhinoceroses (*Diceors bicornis michaeli*;  $n = 5$ ) during pregnancy. Samples were analyzed with EIAs for 20-oxo-pregnanes ( $\circ$  5 $\alpha$ -20-one EIA;  $\bullet$  5 $\beta$ -20-one EIA;  $\square$  4-ene-20-one EIA). Note break of vertical axis.




Fig. 4. Concentrations of immunoreactive 20-oxo-pregnanes in fecal samples of a black rhinoceros (*Diceors bicornis minor*) during pregnancy. Samples were analyzed with EIAs for 20-oxo-pregnanes ( $\circ$  5 $\alpha$ -20-one EIA;  $\bullet$  5 $\beta$ -20-one EIA;  $\square$  4-ene-20-one EIA). Note break of vertical axis.

20-one EIA) were produced. In both assays considerably higher crossreactivities with 20-oxo-pregnanes were obtained, as compared to the 4-ene-20-one EIA. However, the newly developed EIAs had crossreactions with 5 $\alpha$ - and 5 $\beta$ -pregnanes containing a 20-oxo-group and thus were not specific for the 5 $\alpha$ - and 5 $\beta$ -pregnane series, respectively. Therefore, values obtained in these EIAs were comparable, but they were considerably higher than those in the 4-ene-20-one EIA.

In fecal samples of the black rhinoceros "authentic" progesterone, which would have coeluted with  $^3\text{H}$ -progesterone and which had crossreactivities of 100%

in all three EIAs used, was not detectable. HPLC separation showed three major immunoreactive peaks; their elution patterns and the different crossreactivities in the three respective EIAs provided strong evidence that these peaks are 5 $\alpha$ -pregnane-3,20-dione, 5 $\alpha$ -pregnane-3 $\alpha$ -ol-20-one, and 5 $\alpha$ -pregnane-3 $\beta$ -ol-20-one. Although rigorous identification was not achieved (GC/MS was not available), our findings of several 20-oxo-pregnanes are in accordance with other studies in different species. Using progesterone antibodies, the presence of 20-oxo-progesterone metabolites in the feces of primates [Wasser et al., 1991, 1994], felids [Graham et al., 1993; Möstl et al., 1993; Brown et al., 1994; Czekala et al., 1994], ruminants [Desaulniers et al., 1989; Messier et al., 1990; Monfort et al., 1993; Larter et al., 1994], white rhinoceroses [Schwarzenberger et al., 1994, 1995b], and vicunas [Schwarzenberger et al., 1995a] was demonstrated. In addition, radioinfusion studies [Shideler et al., 1993; Brown et al., 1994; Wasser et al., 1994; Palme, personal communication] and studies using HPLC separation with immunoassay analysis [Kirkpatrick et al., 1991; Schwarzenberger et al., 1991, 1992, 1993a,b, 1995a; Heistermann et al., 1993; Shideler et al., 1993; Brown et al., 1994; Wasser et al., 1994] showed that progesterone metabolism and excretion results in several fecal steroid metabolites showing elution patterns similar to that of pregnanes.

The comparisons made in this manuscript deal with progesterone assays and not assays recognizing either a 20 $\alpha$ -OH or a 20 $\beta$ -OH group. Pregnanediol (5 $\beta$ -pregnane-3 $\alpha$ ,20 $\alpha$ -diol) has a 20 $\alpha$ -OH group in position C20 of the steroid molecule, and its crossreactivities in our 20-oxo-assays are <0.1% (see Table 2). However, most pregnanediol antibodies are position-3 conjugates against a 20 $\alpha$ -ol-5 $\beta$ -pregnane and therefore correspond with our definition of group-specific. Pregnanediol antibodies were used for fecal steroid analysis in mares [Kirkpatrick et al., 1991], primates [Heistermann et al., 1993; Shideler et al., 1993], okapis [Schwarzenberger et al., 1993b], and black rhinoceroses [Schwarzenberger et al., 1993a; Berkely, 1994].

Since "authentic" progesterone is not (or to a limited extent only) present in fecal samples, antiserum selection for the analysis of progesterone metabolites is a serious concern. Some of the currently available antibodies for progesterone (namely 11HS- or 6HS-progesterone antisera) are rather specific and show little crossreactivity to other metabolites. However, fecal progesterone metabolites derive from one parent molecule and it is desirable to quantify several metabolites in one assay. In this respect, group-specific assays using steroid C3-conjugate antibodies are superior; they show high crossreactivities with several progesterone metabolites sharing a similar C20 configuration and their measurement is combined in one assay. This concept was also the basis for the "total estrogen assays," which employ 17-hemisuccinate derivatives of estradiol and which recognize the phenolic A ring, common to several naturally occurring estrogens [Niswender et al., 1975].

Instead of using a 5-reduced 20-oxo-pregnane as standard, we used progesterone. Progesterone is easily available, and application-orientated laboratories can use group-specific 20-oxo-pregnane assays for fecal steroid analysis in several animal species without laborious identification of the major species-specific metabolite(s). Since fecal 20-oxo-pregnanes are present in different species, group-specific assays can be used confidently across species. However, assay results will depend upon the summarized crossreactivities of the fecal steroid metabolites. Therefore, absolute values obtained in different assays may vary considerably, so for diagnostic purposes laboratories using different assays might use their own threshold values. Crossreac-

tivities with 20-oxo-pregnanes in the present study were considerably higher than those of a progesterone antibody used in a comparable study [Berkeley, 1994]. Our results were 3–5 times higher, although our calculations were based on wet feces and those of Berkeley [1994] on dry feces. Similarly, a study on feline species using a monoclonal progesterone antibody with high crossreactivities to 5 $\alpha$ -reduced pregnanes reported 10–100-fold higher progesterone metabolite concentrations as compared to other studies [Brown et al., 1994].

The increase in assay specificity enabled us to determine fecal 20-oxo-pregnane values more accurately as compared with the 4-ene-20-one EIA [Schwarzenberger et al., 1993a]. Thus, differences in the fecal pregnane values between nonpregnant and >2 months pregnant animals increased. The increase in accuracy causes an increase in reliability of pregnancy diagnosis. Although this is not of the utmost importance for animals kept in zoos, where several samples can be collected over prolonged time periods, the high assay reliability is important for the study of free-ranging animals. Confirmation of pregnancy from one sample is possible by monitoring the high fecal 20-oxo-pregnane values 3–4 months after mating. These high pregnane values most probably reflect placental steroid production. The increase in fecal steroid excretion in the black rhinoceros after the third month of pregnancy was also described by Berkeley [1994]. Serum hormone levels of >1,000 ng/ml were obtained in a progesterone RIA during the last third of pregnancy [Kock et al., 1991].

During the last quarter of gestation fecal 20-oxo-pregnane values declined. This is in contrast to the pattern of fecal progesterone metabolites containing a 20 $\alpha$ -OH group which, in black rhinoceroses and mares, increased throughout gestation [Schwarzenberger et al., 1991, 1993a]. HPLC separation of fecal immunoreactive progesterone metabolites in black rhinoceroses [Schwarzenberger et al., 1993a] suggested that metabolites with higher polarity became more prominent as pregnancy progressed. A shift in steroid production/metabolism and/or excretion to more hydroxylated and thus more polar progesterone metabolites might explain this phenomenon. The patterns of 5 $\alpha$ -pregnanes containing a 20 $\alpha$ -OH group using group-specific assays need to be elucidated in future investigations. These assays might improve estrous cycle monitoring, since the conversion of radioactively labeled progesterone into 20 $\alpha$ -dihydroprogesterone in a rhinoceros was demonstrated [Hindle and Hodges, 1990], and since urinary analysis of 20 $\alpha$ -dihydroprogesterone was successfully used for estrous cycle monitoring [Hindle et al., 1992].

## CONCLUSIONS

1. Fecal 20-oxo-pregnanes in the black rhinoceros consist of closely related immunoreactive metabolites. Their elution profiles on HPLC and different crossreactivities in the three EIAs used provided strong evidence that these metabolites are 5 $\alpha$ -pregnane-3,20-dione, 5 $\alpha$ -pregnane-3 $\alpha$ -ol-20-one, and 5 $\alpha$ -pregnane-3 $\beta$ -ol-20-one.

2. Values of fecal 20-oxo-pregnanes are determined more accurately using group-specific assays against 5-pregnane-20-one C3 conjugates, as compared to a progesterone antibody.

3. Using the newly developed assays, differences between values of nonpregnant and pregnant animals increased and thus reliability of pregnancy diagnosis was improved.

4. Application of group-specific assays for 20-oxo-pregnanes reaffirms that they might be of general interest for future analysis of fecal progesterone metabolites in several animal species.

## ACKNOWLEDGMENTS

We thank the animal keepers Z. Vondra, P. Petřřílek, M. Kober, J. Ždůvarek, and V. Hák (Zoo Dvůr Kralove); and K. Jahnel, W. Pittermann, S. Petermann, and N. Steinberg (Zoo Frankfurt) for the conscientious collection of the fecal samples; and A. Aichinger, E. Leitner, and A. Kuchar-Schulz for excellent technical assistance.

## REFERENCES

- Adams, G.P.; Plotka, E.D.; Asa, C.S.; Ginther, O.J. Feasibility of characterizing reproductive events in large nondomestic species by transrectal ultrasonic imaging. *ZOO BIOLOGY* 10:247–259, 1991.
- Berkeley, E.V. FECAL STEROID HORMONE ANALYSIS OF OVULATION AND PREGNANCY IN THE BLACK RHINOCEROS (*DICEROS BICORNIS*). M.S. Thesis, Ohio State University, 1994.
- Brown, J.L.; Wasser, S.K.; Wildt, D.E.; Graham L.H. Comparative aspects of steroid hormone metabolism and ovarian activity in felids, measured noninvasively in feces. *BIOLOGY OF REPRODUCTION* 51:776–786, 1994.
- Czekala, N.M.; Durrant, B.S.; Callison, L.; Williams, M.; Millard, S. Fecal steroid hormone analysis as an indicator of reproductive function in the cheetah. *ZOO BIOLOGY* 13:119–128, 1994.
- Czekala, N.M.; Callison, L. Pregnancy diagnosis in the black rhinoceros (*Diceros bicornis*) by salivary hormone analysis. *ZOO BIOLOGY*, in press.
- Desaulniers, D.M.; Goff, A.K.; Betteridge, K.J.; Rowell, J.E.; Flood, P.F. Reproductive hormone concentrations in faeces during the oestrous cycle and pregnancy in cattle (*Bos taurus*) and muskoxen (*Ovibos moschatus*). *CANADIAN JOURNAL OF ZOOLOGY* 67:1148–1154, 1989.
- Graham, L.H.; Raeside, J.I.; Goodrowe, K.L.; Liptrap, R.M. Measurements of faecal oestradiol and progesterone in non-pregnant and pregnant domestic and exotic cats. *JOURNAL OF REPRODUCTION AND FERTILITY (SUPPLEMENT)* 47:119–120, 1993.
- Heistermann, M.; Tari, S.; Hodges J.K. Measurement of faecal steroids for monitoring ovarian function in New World primates, *Callitrichidae*. *JOURNAL OF REPRODUCTION AND FERTILITY* 99:243–251, 1993.
- Hindle, J.E.; Hodges, J.K. Metabolism of oestradiol-17 $\beta$  and progesterone in the white rhinoceros (*Ceratotherium simum simum*). *JOURNAL OF REPRODUCTION AND FERTILITY* 90: 571–580, 1990.
- Hindle, J.E.; Möstl, E.; Hodges, J.K. Measurement of urinary oestrogens and 20 $\alpha$ -dihydroprogesterone during ovarian cycles of black (*Diceros bicornis*) and white (*Ceratotherium simum*) rhinoceroses. *JOURNAL OF REPRODUCTION AND FERTILITY* 94:237–249, 1992.
- Hodges, J.K.; Green, D.I. The development of an enzyme-immunoassay for urinary pregnenediol-3-glucuronide and its application to reproductive assessment in exotic mammals. *JOURNAL OF ZOOLOGY LONDON* 219:89–99, 1989.
- Kasman, L.H.; Ramsay, E.C.; Lasley, B.L. Urinary steroid evaluations to monitor ovarian function in exotic ungulates: III. Estrone sulfate and pregnenediol-3-glucuronide excretion in the Indian rhinoceros (*Rhinoceros unicornis*). *ZOO BIOLOGY* 5:355–361, 1986.
- Kassam, A.A.H.; Lasley, B.L. Estrogen excretory patterns in the Indian rhinoceros (*Rhinoceros unicornis*), determined by simplified urinary analysis. *AMERICAN JOURNAL OF VETERINARY RESEARCH* 42:251–255, 1981.
- Kellie, A.E.; Lichman, K.V.; Samarajeewa, P. Chemistry of steroid-protein conjugate formation. Pp. 33–46 in *STERIOD IMMUNOASSAY, PROCEEDINGS OF THE FIFTH TENOVUS WORKSHOP*. E.H.D. Cameron; S.G. Hillier; K. Griffiths, eds. Cardiff, Wales, UK: Alpha Omega Publishing Ltd., 1975.
- Kirkpatrick, J.F.; Shideler, S.E.; Lasley, B.L.; Turner, J.W. Jr. Pregnancy determination in uncaptured feral horses by means of fecal steroid conjugates. *THERIOGENOLOGY* 35:753–760, 1991.
- Kirkpatrick, J.F.; Gudermuth, D.F.; Flagan, R.L.; McCarthy, J.C.; Lasley, B.L. Remote monitoring of ovulation and pregnancy of Yellowstone bison. *JOURNAL OF WILDLIFE MANAGEMENT* 57:407–412, 1993.
- Klős, H-G.; Frese, R. Foreword to the 5th revised edition of the international studbook of African rhinoceroses. Pp. 12–15 in *INTERNATIONAL STUDBOOK OF AFRICAN RHINOCEROSSES (DICEROS BICORNIS/CERATOTHERIUM SIMUM)*, No. 5. H-G. Klős; R. Frese, eds. Berlin, Germany: Zoologischer Garten Berlin, 1993.
- Kock, N.; Morton, D.; Kock, M. Reproductive parameters in free-ranging female black rhinoceroses (*Diceros bicornis*) in Zimbabwe. *ONDERSTEEPOORT JOURNAL OF VETERINARY RESEARCH* 58:55–57, 1991.
- Kock, R.A.; Wambua, J.; Mwanzia, J.; Saigawa,

- K. Effects of translocation on the black rhinoceros (*Diceros bicornis*)—red and white cell parameters. VERHANDLUNGSBERICHT ERKRANKUNGEN DER ZOOTIERE 37:33–39, 1995.
- Kuckelkorn, B.; Dathe, H.H. Trächtigkeitsdiagnose beim Panzernashorn (*Rhinoceros unicornis*) anhand von Progesteronbestimmungen im Speichel. ZOOLOGISCHER GARTEN N.F. 60: 333–340, 1990.
- Larter, N.C.; Rajamahendran, R.; Sivakumaran, K. Immunoreactive faecal progestins as indicators of reproductive status. VETERINARY RECORD 134:474–475, 1994.
- Messier, F.; Desaulniers, D.M.; Goff, A.K.; Nault, R.; Patenaude, R.; Crete, M. Caribou pregnancy diagnosis from immunoreactive progestins and estrogens excreted in feces. JOURNAL OF WILDLIFE MANAGEMENT 54:279–283, 1990.
- Monfort, S.L.; Schwartz, C.C.; Wasser, S.K. Monitoring reproduction in captive moose using urinary and fecal steroid metabolites. JOURNAL OF WILDLIFE MANAGEMENT 57:400–407, 1993.
- Möstl, E.; Lehmann, H.; Wenzel, U. Gestagens in faeces of mink and cats for monitoring corpus luteum activity. JOURNAL OF REPRODUCTION AND FERTILITY (SUPPLEMENT) 47: 540–541, 1993.
- Niswender, G.D.; Nett, T.M.; Meyer, D.L.; Hagerman, D.D. Factors influencing the specificity of antibodies to steroid hormones. Pp. 61–66 in STEROID IMMUNOASSAY, PROCEEDINGS OF THE FIFTH TENOVUS WORKSHOP. E.H.D. Cameron; S.G. Hillier; K. Griffiths, eds. Cardiff, Wales, UK: Alpha Omega Publishing Ltd., 1975.
- Palme, R.; Möstl E. Biotin–streptavidin enzyme immunoassay for the determination of oestrogens and androgens in boar faeces. Pp. 111–117 in PROCEEDINGS OF THE 5TH SYMPOSIUM ON THE ANALYSIS OF STEROIDS. S. Görög, ed. Budapest, Hungary: Akademiai Kiado, 1993.
- Pryce, C.R.; Schwarzenberger, F.; Döbeli, M. Monitoring fecal samples for estrogen excretion across the ovarian cycle in Goeldi's monkey (*Callimico goeldii*). ZOO BIOLOGY 13:219–230, 1994.
- Ramsay, E.C.; Kasman, L.H.; Lasley, B.L. Urinary steroid evaluations to monitor ovarian function in exotic ungulates: V. Estrogen and pregnanediol-3-glucuronide excretion in the black rhinoceros (*Diceros bicornis*). ZOO BIOLOGY 6:275–282, 1987.
- Ramsay, E.C.; Moran, F.; Roser, J.F.; Lasley, B.L. Urinary steroid evaluations to monitor ovarian function in exotic ungulates: X. Pregnancy diagnosis in perissodactyla. ZOO BIOLOGY 13: 129–147, 1994.
- Schaffer, N.E.; Zainal-Zahari, Z.; Suri, M.S.M.; Jainudeen, M.R.; Jeyendran, R.S. Ultrasonography of the reproductive anatomy in the Sumatran rhinoceros (*Dicerorhinus sumatrensis*). JOURNAL OF ZOO AND WILDLIFE MEDICINE 25: 337–348, 1994.
- Schwarzenberger, F.; Möstl, E.; Bamberg, E.; Pammer, J.; Schmechlik, O. Concentration of progestagens and oestrogens in the faeces of pregnant Lipizzan, Trotter and Thoroughbred mares. JOURNAL OF REPRODUCTION AND FERTILITY (SUPPLEMENT) 44:489–499, 1991.
- Schwarzenberger, F.; Francke, R.; Göldenboth, R. Concentrations of faecal immunoreactive progestagen metabolites during the oestrous cycle and pregnancy in the black rhinoceros (*Diceros bicornis michaeli*). JOURNAL OF REPRODUCTION AND FERTILITY 98:285–291, 1993a.
- Schwarzenberger, F.; Patzl, M.; Francke, R.; Ochs, A.; Buitter, R.; Schaftenaar, W.; DeMeurichy, W. Fecal progestagen evaluations to monitor the estrous cycle and pregnancy in the okapi (*Okapia johnstoni*). ZOO BIOLOGY 12:549–559, 1993b.
- Schwarzenberger, F.; Tomasova, K.; Walzer, C.; Möstl, E. Preliminary results of fecal progestagen evaluations in the white rhinoceros (*Ceratotherium simum*) indicate an estrous cycle length of  $\approx 10$  weeks. BIOLOGY OF REPRODUCTION 50 (SUPPLEMENT) 1:171, 1994.
- Schwarzenberger, F.; Speckbacher, G.; Bamberg, E. Plasma and fecal progestagen evaluations during and after the breeding season of the female vicuna (*Vicuna vicuna*). THERIOGENOLOGY 43:625–634, 1995a.
- Schwarzenberger, F.; Walzer, C. Oestrous cycle induction in a white rhinoceros (*Ceratotherium simum*) monitored by fecal progestagen analysis. VERHANDLUNGSBERICHT ERKRANKUNGEN DER ZOOTIERE 37:79–83, 1995b.
- Shideler, S.E.; Ortuno, A.M.; Moran, F.M.; Moorman, E.A.; Lasley, B.L. Simple extraction and enzyme immunoassays for estrogen and progesterone metabolites in the feces of *Macaca fascicularis* during non-conceptive and conceptive ovarian cycles. BIOLOGY OF REPRODUCTION 48:1290–1298, 1993.
- Wagner, R.A.; Monitoring the reproductive cycle of one southern white rhinoceros. Pp. 14–16 in PROCEEDINGS OF THE ANNUAL MEETING OF THE AMERICAN ASSOCIATION OF ZOO VETERINARIANS. M.S. Silberman; S.D. Silberman, eds. Chicago, Illinois, 1986.
- Wasser, S.K.; Monfort, S.L.; Wildt, D.E. Rapid extraction of faecal steroids for measuring reproductive cyclicity and early pregnancy in free-ranging yellow baboons (*Papio cynocephalus cynocephalus*). JOURNAL OF REPRODUCTION AND FERTILITY 92:415–423, 1991.
- Wasser, S.K.; Monfort, S.L.; Southers, J.; Wildt, D.E. Excretion rates and metabolites of oestradiol and progesterone in baboon (*Papio cynocephalus cynocephalus*) faeces. JOURNAL OF REPRODUCTION AND FERTILITY 101:213–220, 1994.