

RANGING BEHAVIOR AND HABITAT PREFERENCE BY A TRANSLOCATED
POPULATION OF GREATER ONE-HORNED RHINOCEROS (*RHINOCEROS*
UNICORNIS) IN LOWLAND, NEPAL

SHANT R. JNAWALI

Address: King Mahendra Trust for Nature Conservation, Box 3712, Kathmandu, Nepal

Present address: Department of Biology and Nature Conservation, Box 5014, Agricultural
University of Norway, Ås

KEY WORDS: Greater one-horned rhinoceros, *Rhinoceros unicornis*, ranging behavior,
habitat preference, Nepal.

ABSTRACT

Ranging behavior and habitat preferences of a newly translocated population (Bardia) were studied during 1990-1993, and movement pattern was compared with that of the donor population (Chitwan) based on monitoring daytime movement of radiomarked animals. Five years after translocation, the annual home range sizes were $25.1 \pm 9.3 \text{ km}^2$ and $41.8 \pm 4.4 \text{ km}^2$ for adult females ($N = 8$) and males ($N = 2$), respectively ($p < 0.05$). Range sizes of four social groups did not vary between seasons, but during the hot season and the monsoon females with young calves (< 1 year old) had smaller ranges than single females and females with older calves. Seasonal home range size of males varied with social status; when a younger male with a smaller range size displaced an older male with a larger range and became the dominant breeder, his seasonal ranges expanded whereas those of the former dominant's decreased in size.

Annual and seasonal home ranges of animals in the Chitwan donor population were

preferences of this and the donor population have been reported elsewhere (Jnawali, in ms). In another preliminary study, the home range sizes of the Bardia population were found to be much larger compared to the donor population in Chitwan (Wegge and Jnawali 1993). Another purpose of this study was, therefore, to compare the ranging behavior of the translocated founder population with the dense donor population and to examine the reasons for the different spacing pattern observed.

STUDY AREA

The Bardia translocation habitat of about 70 km² is located in the south-western corner of Royal Bardia National Park (81° 20' E and 28° 35' N) (Fig. 1). It consists basically of a narrow strip of alluvial floodplain along Geruwa river. About one third of the area extends between the park boundary and the Indo-Nepalese border in the south. Topographically, the area is flat with elevations between 100 m and 200 m a.s.l. The floodplain is intersected by a large number of small riverlets originating from Geruwa river. The climate, soils, fauna and flora is described earlier (Dinerstein 1979a and 1979b).

The Bardia study area consists of ten distinct habitat types (Table 1). Sal forest, dominated by *Shorea robusta* and *Buchhaniania latifolia*, covers the drier upland in the north-eastern section (Fig. 1). Riverine forest, dominated by *Mallotus philippinensis* and *Syzigium cumini*, grows on poorly drained soils along the riverlets. The floodplain is composed of patches of Tall Grassland dominated by *Saccharum spontaneum*, and successional Khair *Acacia catechu* - Sissoo *Dalbergia sissoo* forest. Phantas (previously cultivated fields) and Wooded Grasslands are dominated mainly by short grasses such as *Imperata cylindrica* (Pokharel 1993). Mixed Hardwood forest occupies a small portion of the study area intermingled with Riverine forest, Khair-Sissoo forest and Wooded Grassland in the northern section of Khauraha island. Bushy pastures are heavily grazed areas along the village boundary outside the park and are mostly located in the southern section. A detailed description of these habitat types is also available elsewhere (Dinerstein (1979a), Jnawali in ms).

In Chitwan, mainly five habitat types are found, viz. Riverine forest, Sal forest, Tall Grassland, Bushy Pasture and cultivated land. The latter four are similar to those found in

reported elsewhere (Jnawali, in ms). The Bardia population were found in Chitwan (Wegge and Jnawali 1993) to compare the ranging behavior of the population and to examine the reasons

located in the south-western corner (Fig. 1). It consists basically of a about one third of the area extends in the south. Topographically, the floodplain is intersected by the Rapti river. The climate, soils, fauna and

habitat types (Table 1). Sal forest covers the drier upland in the north. *Lotus philippinensis* and *Syzigium cumratense* are common in the floodplain. The floodplain is composed of *Stenandrium*, and successional *Khair* (previously cultivated fields) and grasses such as *Imperata cylindrica*. The southern portion of the study area is a Flooded Grassland in the northern grazed areas along the village in the southern section. A detailed description is given by Dinerstein (1979a), Jnawali in

Riverine forest, Sal forest, Tall grassland are similar to those found in

Bardia, but Riverine forest differs in species composition. In Chitwan, this vegetation is richer and dominated by *Trewia nudiflora* with abundant *Bombax ceiba* and *Litsea monopetala*. Besides, two important browse species, *Murraya paniculata* and *Coffea bengalensis*, grow only in Chitwan (Jnawali in ms).

Large tracts of intensively cultivated land and less preferred Sal forest occur outside the boundaries of both study areas. During a two-week period in early winter, several thousand local villagers are permitted to cut grass inside the parks, which, combined with widespread grass burning, reduces the density and cover in the floodplains.

The fauna in both parks is similar, except for nilgai *Boselaphus tragocamelus* and swamp deer *Cervus duvauceli* being confined only to Bardia and gaur *Bos gaurus* only in Chitwan. Large mammals common to both areas besides rhinoceros include tiger *Panthera tigris*, leopard *P. pardus*, sloth bear *Melursus ursinus*, wild boar *Sus scrofa* several species of deer (*Axis axis*, *A. porcinus*, *Cervus unicolor*, and *Muntiacus muntjack* and transient wild elephants *Elephas maximus*.

MATERIALS AND METHODS

The Bardia founder population, consisting of 2 adult males, 5 adult females, 3 subadult females, and 3 subadult males, were captured near Sauraha in Chitwan and translocated and released at the same site in the upper Karnali floodplain, about 10 km north of the Park headquarter (Fig. 1). During 8 years after release in 1986, 9 animals died and 11 calves were born. In late 1994, the population consisted of 15 individuals (3 males, 8 females and 4 calves, estimated at a density of approximately 0.3 animals/km² (Jnawali and Wegge, in ms).

Data from Chitwan used in the present comparison were taken from a subpopulation residing near Sauraha in the northeast part of the park. This population has been intensively studied by Laurie (1978 and 1982) and Dinerstein and Price (1991), and consists of > 100 animals in prime habitats along the Rapti river, estimated at a density of 8-10 animals/km² (Dinerstein and Price 1991).

In 1990, nine of the 10 founders then alive in Bardia were captured and equipped with radio transmitters in the 142 Mhz frequency range. In addition, one subadult female born in Bardia was also radio instrumented in 1992. All animals were immobilized using the same

drugs and techniques as described by Dinerstein et al. (1990). The drug dose was increased to 2.5 - 3 mg of M99 and 1.5 mg of acepromazine to ensure rapid immobilization. Animals were tracked during day-time, using a portable receiver and two element antennas, and located by triangulation, cross bearings or by direct approach. Tracking was carried out by elephants during the monsoon and by car and bicycle during the dry season. On average, all animals were tracked 3 times per week. The tracking periods were divided into three seasons - winter season (November-February), hot season (March-June) and Monsoon (July - October).

In Bardia, a total of 1641 locations from 10 rhinoceros were obtained between February 1990 and March 1993. In Chitwan, data (N = 652) were obtained from five animals (1 male and 4 females) tracked during 1985-87. Locations were plotted on aerial photos (1:35000 in Bardia and 1:30000 in Chitwan) and later transferred to topographic maps. In both study areas, composition of vegetation types were identified and mapped based on transect sampling and later measured by a digital planimeter (Jnawali, in ms).

Both seasonal and annual home ranges were drawn as convex polygons using the "modified minimum area method" (Harvey and Barbour 1965) as modified by Wegge and Larsen (1987): the distance between the two widest locations was divided in half. A line was then drawn clockwise between all successive outermost points that were spaced shorter than this half maximum distance. Locations further away than the maximum distance were defined as excursions and were excluded in the estimate of home range sizes. The size of a home range is a function of sample size (Voigt and Tinline 1980). In the present study an asymptotic relationship between home range size and number of location was obtained after 28 day locations. Hence, animals with less than 28 locations were excluded from the analysis of seasonal home ranges.

In Bardia, > 95% of the consecutive locations of each individual animal were spaced at intervals of 48 hours or more. Thus, autocorrelation was not considered a problem with the Bardia data set. In Chitwan, animals were tracked more frequently. To avoid autocorrelation and make data comparable with Bardia, only locations between 48 and 51 hours intervals were used.

The occupancy center (OC), or geographical center of a home range, was defined as the intersection of the average distance of locations measured along the x and y coordinates (Wegge and Larsen 1987). 50% of the day locations nearest to the occupancy center were used to estimate the core area of each seasonal home range. The size of home range and core

1990). The drug dose was increased
are rapid immobilization. Animals
two element antennas, and located
tracking was carried out by elephants
y season. On average, all animals
divided into three seasons - winter
d Monsoon (July - October).

inceros were obtained between
) were obtained from five animals
ns were plotted on aerial photos
nsferred to topographic maps. In
identified and mapped based on
er (nawali, in ms).

n as convex polygons using the
965) as modified by Wegge and
s was divided in half. A line was
nts that were spaced shorter than
maximum distance were defined
ange sizes. The size of a home
1989). In the present study an
er of location was obtained after
were excluded from the analysis

individual animal were spaced
t considered a problem with the
ently. To avoid autocorrelation
een 48 and 51 hours intervals

a home range, was defined as
along the x and y coordinates
to the occupancy center were
the size of home range and core

areas was measured by a digital planimeter. Home range stability was determined
calculating overlap of successive seasonal home ranges and core areas. Range length, defined
as the minimum distance moved by an animal during a given time interval, was estimated
using the method described by Wegge (1985). In this study, an interval of 50 hrs (3 days)
used to estimate and compare movement pattern within home ranges.

Habitat preferences were examined at three levels: (a) comparing frequency
distributions of animal location on habitat types with distribution of habitats within the total
study area, (b) comparing frequency distribution of habitats within seasonal core areas with
frequency distribution of habitats within total study area, and (c) comparing frequency
distribution of animal locations on habitat types with frequency distribution of habitats within
seasonal home ranges of each individual animal. For the latter, each animal's frequency
distribution on each of the three main habitat types, viz. Tall Grassland, Khair-Sissoo, and
Riverine forest, was plotted against the frequency of the same habitat within the animal's
home range. Plots on the 45° median line indicated that the animal made no active selection
for that particular habitat (Kolstad et al. 1985).

Preference for particular habitats was determined by Bonferroni confidence intervals
(Neu et al. 1974, Byers et al. 1984). Ivlev's Electivity Index (IEI) was employed to estimate
the degree of preference or avoidance of each habitat types (Ivlev 1961). Seasonal habitat
preferences by individual animals were determined by the relative proportion of individual
animal "plots" which fell on each side of the median line and were analyzed by the Wilcoxon
matched pair signed-rank test.

RESULTS

Home range

Average annual home range sizes of females and males of the translocated population
in Bardia were estimated at $25.1 \pm 9.3 \text{ km}^2$ and $41.8 \pm 4.4 \text{ km}^2$, respectively, whereas in Chitwan
they were only 2.9 km^2 (females) and 3.3 km^2 (males) (Table 2). Similarly, seasonal home
ranges of females and one male in Chitwan were < 10% that of Bardia animals. The
corresponding core areas of Bardia animals were also considerably larger than those of

Chitwan animals.

In Bardia, annual home range size of females with calves were not different from that of single females (Wilcoxon rank-sum test, $W_1 = 8$, $n_1 = 2$, $n_2 = 6$; $p > 0.05$). The annual home range size of males was significantly (Wilcoxon rank-sum test, $W_1 = 19$, $n_1 = 2$, $n_2 = 8$; $p < 0.05$) larger than the average annual home range size of single females and females with calves lumped together (Table 3).

In Bardia, no statistical differences were detected between size of seasonal home ranges of four different social groups (Kruskal-Wallis test, $p > 0.05$, all groups) (Table 4). In the hot and monsoon seasons, the size of home ranges of females with older calves (> 1 year) and single females were both significantly larger than the females with small calves (< 1 year) (Wilcoxon rank-sum test, $p < 0.05$, all), but no such differences were observed in winter. In all three seasons, mean home range sizes of males were larger than that of females with big calves and single females combined, but none of them were statistically significant (Wilcoxon rank-sum test, $p > 0.05$, all). Among the two males, size of home ranges changed with social status (Table 4). When the younger, subadult male (337) replaced the older, dominant breeder (275) in the monsoon, home range sizes of both changed in opposite directions.

Females tracked in consecutive seasons (hot = 5, monsoon = 4 and winter = 2) were used to detect changes in home range size from one year to the next. In general, ranges increased slightly (8.6%) in consecutive seasons, but none of the changes were statistically significant (Wilcoxon matched pair signed-rank test, $p > 0.05$, all) (Table 5).

All instrumented animals remained mainly within a narrow strip of the floodplain along Geruwa river with remarkable overlap in seasonal home ranges (Table 6). For females, the mean seasonal home range overlap ranged from 54.9% to 74.8%, whereas for males it ranged from 41.2% to 51.3%. The seasonal core area overlap ranged from 21.6% to 43.8%

(females) and from 18.6% to 21.6% (males). Overlap was largest during the hot/monsoon, smallest during monsoon/winter seasons. Seasonal overlap of males was consistently smaller than that of females (Table 6).

In Bardia all radio marked animals shifted their occupancy centers between seasons (Fig. 2), in spite of large seasonal overlap in home ranges (Table 6). For males the mean distances between OCs ranged from 3.3 km to 6.2 km, whereas for females they ranged from 1.1 km to 3.1 km. The difference between sexes were significant in the winter/hot and hot/monsoon seasons (Wilcoxon rank sum test, $p < 0.05$, both) but not between the monsoon and winter (Wilcoxon rank sum test, $W_1 = 13$, $n_1 = 2$, $n_2 = 8$, $p > 0.05$).

In Bardia, animals moved extensively within their home ranges (Fig. 3). Mean linear distance within ca 50 hours intervals ranged from 2.3 km to 4.1 km for males and 2.5 to 2.9 km for females (Table 6). Thus, within few days animals traversed most of their home ranges. For example, one female made a journey of about 11 km (location 23 to 24) in < 35 hours (Fig. 3). On average, males tended to move longer than females, but the difference was only significant in the hot season (Wilcoxon rank sum test, $W_1 = 13$, $n_1 = 2$, $n_2 = 5$, $p < 0.05$).

With smaller annual and seasonal home ranges in Chitwan, distances between seasonal OCs and range lengths were also much smaller there than in Bardia (Table 6). However, movement in relation to home range size was almost as large; animals moved less but covered nearly the same proportion of their home range within the same time interval. Compared with Bardia, extent of seasonal home range overlap was larger among Chitwan females, except for the hot/monsoon season. The male here had larger overlaps in all three seasons.

Habitat use

Bardia animals preferred Khair-Sissoo forest, Riverine forest and Tall Grassland and avoided Sal forest, Wooded Grassland, and River and Riverbeds (Table 7). Mixed Hardwood forest was used according to availability. Among preferred habitat types, highest preference was observed for Riverine forest (IEI = 0.44). Sal forest was the most avoided (IEI = - 0.93), and animals were not located in Bushy pastures, Phanta and Agricultural lands during daytime.

Riverine forest was preferred in all three seasons by females (Table 8 and Fig. 4), whereas males preferred this habitat only during the hot and winter seasons (Table 8). When seasonal use of habitats was tested on an individual animal level, this habitat was used significantly more in the hot and winter seasons (Wilcoxon matched pair signed-rank test, $t = 55$ and 54 , respectively, $n = 10$, $p < 0.01$, both) (Fig. 5).

Khair-Sissoo forest was also preferred in all three seasons by females (Table 8, Fig. 4), whereas males preferred this habitat type only during winter. Test for seasonal use by individual animals confirmed that this type was preferentially used mainly during winter (Wilcoxon matched pair signed-rank test $t = 51$, $n = 10$, $p < 0.01$) (Fig. 5). This was also reflected in the composition of habitat types in the seasonal core areas. Highest proportion ($\bar{x} = 40.4 \pm 5.4SD$) of Khair-Sissoo forest was recorded during winter (Table 9).

Tall Grassland was preferred during hot and monsoon seasons by females whereas males preferred this habitat type during the monsoon only (Table 8). Both sexes used very little of this habitat during winter (Table 8). Test for seasonal use by individual animals also disclosed that this habitat was preferentially used during the monsoon and hot seasons (Wilcoxon matched pair signed-rank test, $p < 0.01$) (Fig. 5). Furthermore, core areas contained the highest proportion of this habitat during the monsoon ($\bar{x} = 32.3 \pm 2.9SD$), confirming that

Tall Grassland was especially important during this season (Table 9).

There were some notable differences between the habitat use of males and females (Fig. 6). Most pronounced differences were observed in the hot season. Females used significantly more Khair-Sissoo and Tall Grassland than males, whereas males used more Riverine forest than females (Chi Square tests, $p < 0.05$, all). In winter, Khair-Sissoo was much preferred by both sexes with no differences between males and females. However, Riverine forest was then used more by males. In the monsoon, Tall Grassland was much preferred by both sexes, but with differential usage between males and females. In this season Riverine forest was used more by females ($\chi^2 = 9.37$, $p < 0.05$).

DISCUSSION

Ranging pattern

Five years after translocation to Bardia, the average annual home range of individual animals was estimated at about 28.5 km². The annual home range size of animals in the Chitwan donor population was 3.0 km², or only 1/9 of that. In Chitwan, radio instrumented animals basically remained in the prime habitat of Icharni island. However, occasionally they excused across the Rapti river in the west and to Bushy pastures and cultivation in the northeast. These areas are not included in the home ranges in the present study. Their inclusion would increase the home range size by a maximum of 20-30 percent, which still make the Chitwan home ranges much smaller than in Bardia. Laurie (1982) also estimated a smaller mean home range size of 4.3 km² for Icharni animals.

The discrepancies between home range size in Bardia and in Chitwan may be due to

non-settled ranging behavior, lower animal density or poorer habitat quality in Bardia.

Because data on ranging behavior was collected only five years after translocation, animals in Bardia may not yet have settled down. However, seasonal home ranges did not decrease from 1990/91 to 1991/92. In fact, they tended to increase (Table 5), although large individual differences were observed. Seasonal ranges were also spatially quite stable from one year to the next, and supplementary data collected in 1993 supported the general pattern that individual animals continued to roam over the same large areas in which they had settled in 1990.

Kiwia (1989) and Pienaar et al. (1993) found an inverse relation between the home range size and density of black *Diceros bicornis* and white *Ceratotherium simum simum* rhinoceros, respectively. In high density areas, movement of individual animals may be restricted by animals in neighbouring areas, as found among African buffaloes *Syncerus caffer* (Funston et al. 1994). Here, the borders of home ranges of individual herds were dictated mainly by the proximity of neighbouring herds. But rhinoceros are basically non-territorial and solitary animals (Laurie 1978). Hence, the size of home ranges is not expected to be influenced by the presence of other conspecifics.

However, the extremely low animal density in Bardia (ca 1 animal per 5 km²) may have triggered extensive movement unrelated to habitat quality per se. The highly skewed sex ratio of only one mature male/8 breeding age females and a widely spaced distribution pattern of females may have induced movement by both sexes searching for mates. Indirect evidence of ovulations not leading to conception indicates that females, on occasion, did not make contact with the dominant male when in estrous (Jnawali and Wegge in ms). The pronounced increase in movement by the younger male after he became the dominant breeder also suggests that mating behavior may have affected space use.

In general, home range size is likely to increase when large numbers of animals exploit the same resources (Dunham 1981). Thus, assuming a similar resource base in both areas, home range of Chitwan animals is expected to be larger, and not smaller, than those of Bardia animals. Based on resource requirements, home ranges of rhinoceros are probably determined by productivity, availability and distribution of seasonal foods and access to habitats for shelter, heat avoidance and wallowing. Tall Grasslands and Riverine forests are the critical habitats for the greater one-horned rhinoceros (Laurie 1978, Dinerstein and Price 1991, and this study). Besides, agricultural crops become an important component of their diet late in the monsoon and during the early part of the dry season (Laurie 1978, Jnawali 1989). In Chitwan, continuous blocks of Riverine forest surrounded by large tracts of grassland combined with closely located agriculture reduce animals movement (Fig. 7). Unlike in Chitwan, in Bardia the preferred habitats are fragmented in small patches unevenly distributed throughout the north-south elongated floodplain (Fig. 7). This probably leads not only to larger shifts in location of seasonal occupancy centers but also to more extensive movements within each season. Hitchens (1969) and Larter and Gates (1993) also observed larger home ranges of black rhinoceros and wood bison *Bison bison*, respectively, in areas where food resources were more dispersed, and Owen-Smith (1973) reported that white rhinoceros extended their ranges during winter when water was in short supply.

Moe and Wegge (1995) stated that the fine grained habitat pattern in Bardia compared to Chitwan reduced the home range size of axis deer *Axis axis*, in spite of a much higher animal density there. But habitats preferred by rhinoceros differ from those preferred by axis deer. The daily food intake by rhinoceros of about 15-20 kg of organic matter (Foose 1982, cited in Owen-Smith 1988) is extremely high compared to small ungulates. In Bardia, individual patches of preferred habitat are probably too small to supply sufficient quantity of

high quality food for an extended period of time. As a result, animals need to move from one patch to the next in order to meet their high daily food requirement.

However, habitat quality may also be poorer in Bardia. The productivity of the staple food plant - *Saccharum spontaneum* - is probably lower than in Chitwan. Although no productivity study has yet been made in Bardia, indirect evidence indicates that this may be the case. Compared to Chitwan the floodplain in Bardia is drier, probably leading to less vigorous sprouting of *Saccharum* during the dry season, with animals exploiting proportionally less grasses and more browse than Chitwan animals during this time (Jnawali in ms). This may explain the larger home ranges in Bardia during the dry season, but not in the monsoon.

All seasonal home ranges of Bardia animals were larger than those of the donor population in Chitwan. Tall Grassland was the most preferred habitat type during the monsoon. In Bardia, this habitat is scattered in small patches ($\bar{x} = 10.4$ ha) along the river and riverlets, spaced on average 0.99 km apart, in contrast to Chitwan where Tall Grasslands occur in large blocks (Fig. 7). More extensive movement during the monsoon by Bardia animals may therefore be explained by smaller patch size and lower relative proportion of this preferred habitat type in Bardia, rather than inferior quality during this period.

Riverine forest was preferred during the hot season in Bardia. Like with Tall Grassland, Riverine forest is also dispersed in fragmented patches (mean 35 ha) about 2.7 km apart compared with Chitwan where patches are much larger. Furthermore, Riverine forests in Chitwan are richer in preferred food plants (Jnawali in ms.), which probably also contributes to more extensive movement and larger home ranges during the hot season in Bardia.

Less access to cultivated lands probably also increases the home range of Bardia

animals
habitats
river in
lands oc
winter w
L
males w
mainly be
the seaso
movemen
explained
adult male
On
other fema
of young c
Habitat use
In B
(Fig. 6). *Ma*
type, of wh
tenuis, a cli
mainly durin
bushes of C
shelter durin

is need to move from one
productivity of the staple
in Chitwan. Although no
indicates that this may be
probably leading to less
with animals exploiting
during this time (Jnawali
the dry season, but not in
than those of the donor
habitat type during the
(0.4 ha) along the river and
an where Tall Grasslands
the monsoon by Bardia
relative proportion of this
this period.
Bardia. Like with Tall
mean 35 ha) about 2.7 km
thermore, Riverine forests
s), which probably also
during the hot season in
the home range of Bardia

animals. The Sauraha donor population occupies a concentration of naturally highly preferred habitats close to cultivated lands. In Bardia, access to agriculture is limited by the Geruwa river in the west and large tracts of less preferred Sal forest in the east. Accessible cultivated lands occur mostly in the southern section which triggers animal movement southwards during winter when crop raiding becomes prevalent.

Like in black rhinoceros (Kiwia 1989), average annual home range size of Bardia males was significantly larger than the average annual home range of females. This was mainly because their seasonal home ranges were more spaced than those of females. However, the seasonal home ranges of males were also larger, probably due to males' extensive movement in search of females. Furthermore, larger home ranges among males may also be explained by differences in areas needed to fulfill metabolic requirements (McNab 1963), as adult males are 20-25% larger than females (Laurie 1982).

Only during the monsoon was home range size of females with calves smaller than other female social groups. The flooded riverlets may have limited more extensive movement of young calves during this season.

Habitat use

In Bardia, Riverine forest was heavily utilized during winter and hot seasons (Fig. 6). *Mallotus philippinensis* and *Syzigium cumini* are dominant tree species in this habitat type, of which *Mallotus* constitutes an important part of the diet (Jnawali in ms). *Calamus tenuis*, a climbing palm also basically confined to the Riverine forest, was also exploited mainly during the hot and winter seasons. Furthermore, Riverine forest with unpenetrable bushes of *Calamus*, particularly in the southern section of the study area, also provides good shelter during winter when animals raid crops during night-time.

Khair-Sissoo forest was also preferentially used during winter. With a dense and abundant understorey vegetation, consisting mainly of *Murraya koinigii*, *Pogostemon bengalensis* and *Callicarpa macrophylla*, this habitat type provides suitable shelter when the Tall Grassland becomes more open after grass cutting and burning in early winter. Furthermore, *Callicarpa* is an important food plant during this season. Females preferred Khair-Sissoo forest also during the hot season and the monsoon. During the hot season animals voraciously exploit *Dalbergia sissoo* and *Callicarpa macrophylla* (Jnawali in ms). Monsoon use of Khair-Sissoo forest was probably related to shelter to escape from blood sucking tabanid flies *Tabanus sp.*, which are prevalent in Tall Grassland during that part of the year.

Tall Grassland in the floodplain was highly preferred by both male and females during the monsoon, when animals mainly exploit grasses common in this habitat. Preferred *Saccharum spontaneum*, *S. bengalensis*, *Arundo donax* and *Erianthus ravennae* make up about 70% of the diet in this season (Jnawali in ms). Tall Grassland also attracts animals during the hot season when new flush becomes available after the first rains late in the hot season.

Other habitats in Bardia - Sal forest, Mixed Hardwood forest and Wooded Grassland - received very little use. These habitats are of little importance in terms of food plants. Common grasses in the latter two habitats (*Imperata cylindrica* and *Desmostachia bipinnata*) receive some use only during the early sprouting stages, and sal seedlings were not eaten (Jnawali, in ms). Also, with an open understorey vegetation neither of these habitats provide good shelter. River and riverbeds were used relatively more during the hot season when animals wallowed frequently to escape from the heat. Animals were never located in Phanta and Bushy pastures during daytime. Phantas are devoid of both preferred food plants and good shelter. No use of Bushy pastures during the day was mainly due to disturbances, as all

Bushy p
I
animals
more ext
to low
Although
larger sp
Unlike in
and distri
addition,
agricultur
Riverine I
reduces an

I w
permission
logistic sup
Smithsonian
Cooperation
the field wo
and Man B.

ter. With a dense and
at *koinigii*, *Pogostemon*
suitable shelter when the
urning in early winter.
ason. Females preferred

During the hot season
phylla (Jnawali in ms).
er to escape from blood
sland during that part of

male and females during
this habitat. Preferred
ravennae make up about
tracts animals during the
ate in the hot season.

and Wooded Grassland -
n terms of food plants.

Desmostachia bippinata)
seedlings were not eaten
of these habitats provide
ng the hot season when
e never located in Phanta
preferred food plants and
due to disturbances, as all

Bushy pastures lie outside the park boundary and are used for livestock grazing.

Five years after release in Bardia, spacing behavior showed that the translocated animals had settled in established home ranges. Large size of annual and seasonal ranges and more extensive movement compared with the donor population in Chitwan were mainly due to low animal density combined with different spatial distribution of preferred habitats. Although Tall Grassland may be less productive in Bardia, this is unlikely to explain the larger space use there, because animal density was too low to induce competition for food. Unlike in Chitwan, in Bardia the preferred Riverine forest and Tall Grasslands are fragmented and distributed in small patches which contribute to more extensive seasonal movements. In addition, the huge Geruwa river and large tracts of unattractive Sal forest limit access to agricultural crops, leading animals to move southwards during winter. In Chitwan, large tracts Riverine forest surrounded by Tall Grassland combined with nearby agriculture probably reduces animals' movement to obtain seasonal foods.

ACKNOWLEDGEMENTS

I wish to thank the Department of National Park and Wildlife Conservation for permission to carry out this work and King Mahendra Trust for Nature Conservation for logistic support. Data from Chitwan were collected during a five year project funded by the Smithsonian Institute directed by E. Dinerstein. The Norwegian Agency for Development Cooperation funded the field work in Bardia. P. Wegge provided continuous help throughout the field work in Bardia, and with data analyses and write up of this manuscript. Gagan Singh and Man B. Lama were helpful during animal capture and radio-tracking in Bardia.

REFERENCES

- Anstey, D. 1987. Report on stage two of the reintroduction of the greater one horned rhinoceros to th0e Royal Bardia Wildlife Reserve, Nepal. report, DNPWC/KMTNC. 12 pp.
- Bhattacharya, A. 1993. The status of the Kaziranga rhino population. Tiger Paper 20:1-6.
- Bauer J.J. 1988. A priliminary assessment of the reintroduction success of the Asian one-horned rhinoceros (*Rhinoceros unicornis*) in Bardia wildlife reserve, Nepal. Tiger Paper, October-December:26-32.
- Byers, R.C., Steinhorst, R.K., and Krausman, P.R. 1984. Clarification of a technique for analysis of utilization-availability data. J. Wildl. Manage. 48:1050-1053.
- Caughley, G. 1969. Wildlife and recreation in the Trisuli Watershed and other areas in Nepal. HMG/FAO/UNDP Trisuli Watershed and Development Project. Project Report No. 6. Kathmandu. 54pp.
- Damuth, J. 1981. Home range, home mage overlap, and species energy use among herbivorous mammals. Biol. J. Linn. Soc. 15:185-193.
- Dinerstein, E. 1979a. An ecological survey of Royal Karnali-Bardia Wildlife Reserve, Nepal. Part I: Vegetation modifying factors, and successional relationships. Biol. Conserv. 15:127-149.
- Dinerstein, E. 1979b. An ecological survey of the Royal Karnali-Bardia wildlife reserve, Nepal. Part II: Habitat/animal interactions. Biol. Conserv. 18: 5-38.
- Dinerstein, E., and Price, L. 1991. Demography and habitat use by a Greater One-horned rhinoceros in Nepal. J. Wildl. Manage. 55:401-411.

- Dinerstein, E., Shrestha, S., and Mishra, H.R. 1990. Capture, chemical immobilization, and radio-collar life for greater one-horned rhinoceros. *Wildl. Soc. Bull.* 18:36-41.
- Foose, T.J. 1982. Trophic Strategies of ruminant vs nonruminant ungulates. Ph.D. dissertation. University of Chicago.
- Funston, P.J., Skinner, J.D., and Dott, H.M. 1994. Seasonal variation in movement patterns, home range and habitat selection of buffaloes in a semi-arid habitat. *Afr. J. Ecol.* 32:100-114.
- Gurung, K.K. 1989. The Indian rhinoceros. Pp 87-89. In: *Indian wildlife* (Israel and Sinclair eds.). APA Publications, Singapur. 380pp.
- Harvey, M.J., and Barbour, R.W. 1965. Home range of *Microtus ochrogaster* as determined by a modified minimum area method. *J. Mammology* 46:398-402.
- Hitchins, P.M. 1969. The influence of vegetation types on sizes of home ranges of black rhinoceros in Hluhluwe game reserve, Zululand. *The Lammergeyer* 10:81-85.
- Ivlev, V.S. 1961. *Experimental ecology of the feeding fishes*. Yale University Press, New Haven, Conn.
- Jnawali, S.R. 1989. Park-people conflict: Interaction between greater one-horned rhinoceros and people adjacent to the Royal Chitwan National Park in Sauhara area. M.Sc. thesis. Agri. Univ. of Norway. 86pp.
- Jnawali, S.R., and Wegge, P. 1993. Space and habitat use by a small re-introduced population of greater one-horned rhinoceros (*Rhinoceros unicornis*) in Royal Bardia National Park in Nepal - A preliminary report. Pp. 208-217. In: *Rhinoceros Biology and Conservation* (O.A. Ryder, ed.), Proceeding of an International Conference, Zool. Society, San Diego, USA.

- Jnawali, S.R., and Wegge, P. 1991. Is there room for endangered large mammals in a developing country? Preliminary results from a field study on the conservation of greater one-horned rhinoceros (*Rhinoceros unicornis*) in Royal Bardia National Park in Nepal. *Faginfo* 123:145-160.
- Khan, M.K. bin., and Foose, T.J. 1994. Chairman's report: Asian rhino specialist group. *Pachyderm* 18:3-8.
- Kiwia, H.Y.D. 1989. Ranging patterns of the black rhinoceros (*Diceros bicornis*) in Ngorongoro Crater, Tanzania. *Afr. J. Ecol.* 27:305-312.
- Kolstad, M., Bø, T., and Wegge, P. 1985. The habitat and ecology of black grouse (*Tetrao tetrix* L.) during spring and summer in east Norway. *Medd. Norsk Viltforsk.* 13:1-46.
- Larter, N.C., and Gates, C.C. 1994. Home-range size of wood bison: Effects of age, sex, and forage availability. *J. Mammalogy* 75:142-149.
- Laurie, W.A. 1978. The ecology and behavior of the greater one-horned rhinoceros. Ph.D. dissertation. Cambridge University. 450pp.
- Laurie, W.A. 1982. Behavioral ecology of the Greater One-horned rhinoceros (*Rhinoceros unicornis*). *J. Zool., Lond.* 196: 307-341.
- McNab, B.K. 1963. Bioenergetics and the determination of home range size. *Am. Nat.* 97:133-140.
- Mishra, H.R. 1982. Balancing human needs and conservation in Nepal's Royal Chitwan National Park. *Ambio* 11:246-251.
- Mishra, H.R., and Dinerstein, E. 1987. New zip codes for resident rhinos in Nepal. *Smithsonian Magazine* 18:67-73.
- Moe, S.R., and Wegge, P. 1995. Spacing behaviour and habitat use of axis deer (*Axis axis*) in lowland Nepal. *Can. J. Zoology*. In Press.

of large mammals in a
on the conservation of
of Bardia National Park
rhino specialist group.
(*Diceros bicornis*) in
of black grouse (*Tetrao*
orsk Viltforsk. 13:1-46.
Effects of age, sex, and
orned rhinoceros. Ph.D.
rhinoceros (*Rhinoceros*
e range size. Am. Nat.
epal's Royal Chitwan
resident rhinos in Nepal.
e of axis deer (*Axis*

- Nepal, S.K., and Weber, K.E. 1993. Struggle for existence: Park-people conflict in the Royal Chitwan National Park, Nepal. Asian Institute of Technology, Thailand. 199pp.
- Neu, C.W., Bayers, C.R., and Peek, J.M. 1974. A technique for analysis of utilization-availability data. *J. Wildl. Manage.* 38:541-545.
- Owen-Smith, N. 1973. The behavioural ecology of the white rhinoceros. Ph.D. dissertation. University of Wisconsin.
- Owen-Smith, N. 1988. Megaherbivores: Influence of very large body size on ecology. Cambridge Univ. Press, New York. 396pp.
- Pelinck, E., and Upreti, B.N. 1972. A census of rhinoceros in Chitwan National Park and Tamaspur forest, Nepal. HMG/FAO/UNDP National Park and Wildlife Conservation Project, Internal Report, Kathmandu. 10pp.
- Pienaar, D.J., Bothma, J.D.P., and Theron, G.K. 1993. White rhinoceros range size in the south-western Kruger National Park. *J. Zool., Lond.* 229:641-649.
- Pokharel, S.K. 1993. Floristic composition, biomass production, and biomass harvest in the grassland of the royal Bardia National Park, Bardia, Nepal. M.Sc. thesis, Agricultural University of Norway, Ås, Norway. 66pp.
- Sale, J.B. 1986. Rhinos re-established in Uttar Pradesh. *Indian Forester*, October:945-948.
- Sale, J.B., and Singh, S. 1987. Reintroduction of greater Indian rhinoceros into Dudhwa National Park. *Oryx* 21:81-84.
- Sharma, U.R., 1991. Park people interactions in Royal Chitwan National park, Nepal. Ph.D. dissertation. Univ. of Arizona. 274pp.

- Sinha, S.P., and Sawarkar, V. B. 1993. Management of the reintroduction greater one-horned rhinoceros (*Rhinoceros unicornis*) in Dudhwa National Park Uttar Pradesh, India. Pp. 218-227. In: Rhinoceros Biology and Conservation (O.A. Ryder, ed.). Proceeding of an International conference, Zool. Society, San Diego, USA.
- Vigne, L., and Martin, E.B. 1994. The greater one-horned rhino of Assam threatened by poachers. *Pachyderm* 18:28-43.
- Voigt, D.R., and Tinline, R.R. 1980. Strategy for radio tracking data. Pp. 387-404. In: A hand in biotelemetry and radio-tracking, C.J. Amlaner and D.W. McDonald (eds.). Pergamon Press Ltd., Oxford.
- Wegge P. 1985. Spacing pattern and habitat use of capercaillie hens in spring. *Int. Symp on Grouse* 3:499-513.
- Wegge, P., and Larsen, B.B. 1987. Spacing of adult and subadult male Common Capercaillie during the breeding season. *The Auk* 104:481-490.
- Wegge, P., Inawali, S.R., and Moe, S.R. 1990. Bardia Conservation Research Program: Progress report 1990. Department of Biology and Nature Conservation, Agricultural University of Norway, King Mahendra Trust for Nature Conservation, Nepal, and Department of National Parks and Wildlife Conservation, Nepal. 37pp.

Table 1
 Habitat
 Sal fore
 Riverine
 Khair-Si
 Mixed f
 forest
 Tall Gra
 Wooded
 Phanta
 Bushy Pa
 River and
 Cultivated

ction greater one-horned
 Uttar Pradesh, India. Pp.

der, ed.). Proceeding of

of Assam threatened by

a. Pp. 387-404. In: A

D.W. McDonald (eds.).

in spring. Int. Symp on

le Common Capercaillie

n Research Program:

onservation, Agricultural

Conservation, Nepal, and

Nepal. 37pp.

Table 1. Size (km²) and description of the habitat types of the study area in Bardia.

Habitat type	Size	Percentage of study area	Broad description
Sal forest	7.68	11.2	Upper dry land dominated by <i>Shorea robusta</i> and <i>Buchana latifolia</i> .
Riverine forest	6.81	9.9	Poorely drained mixed forest dominated by <i>Mallotus philippinensis</i> , <i>Syzigium cumini</i> and <i>Ehretia laevis</i> .
Khair-Sissoo forest	13.54	19.6	<i>Acacia catechu</i> and <i>Dalbergia sissoo</i> dominated forest growing along the old riverbeds
Mixed Hardwood forest	0.98	1.5	<i>Oegenia dalbergioides</i> and <i>Schleichera trijuga</i> intermixed forest
Tall Grassland	8.19	11.9	Tall grassland communities on the floodplain dominated by <i>Saccharum spontaneum</i> and <i>S. bengalensis</i> . <i>Arundo donax</i> dominates the banks of riverlets.
Wooded Grassland	3.13	4.5	<i>Imperata cylindrica</i> dominated grassland with sparsely distributed tree species, e.g. <i>Bombax ceiba</i> , <i>Bahunia racemosa</i> and <i>Melastoma philippinensis</i> .
Phanta	0.95	1.4	Old cultivated open dry fields dominated by <i>Imperata cylindrica</i> .
Bushy Pasture	4.80	6.9	Heavily grazed meadows dominated by <i>Imperata cylindrica</i> , <i>Chrysopogon aciculatus</i> and <i>Cyanodon dactylon</i> with scattered bushes of <i>Ziziphus mauritiana</i> and <i>Callicarpa macrophylla</i> .
River and riverbed	21.4	31.1	River, riverlets and seasonally flooded bare river beds covered with sand and boulders.
Cultivated land	1.4	2.0	Seasonally irrigated paddy fields and dry corn fields.

Table 2. Seasonal and annual home ranges (HR) and core areas (CA) of Bardia and Chitwan animals (in km²±SD) in 1990-1993.

	Females				Males*			
	Bardia		Chitwan		Bardia		Chitwan	
	HR	CA	HR	CA	HR	CA	HR	CA
Winter	15.6±4.5	2.9±1.7	1.2±0.4	0.2±0.1	20.7±10.3	2.6±0.6	1.1	0.1
Hot	13.3±4.9	3.1±1.4	1.1±0.4	0.2±0.1	21.2±11.1	3.7±0.9	2.0	0.6
Monsoon	14.4±7.1	2.7±1.0	1.3±0.5	0.3±0.1	14.4± 7.1	2.9±0.2	1.4	0.4
All year	25.1±9.3	8.5±2.5	2.9±0.9	0.6±0.2	41.8± 4.4	9.3±3.9	3.3	0.9

* Only 2 males in Bardia and 1 male in Chitwan

of Bardia and Chitwan

Males*	Chitwan	
	HR	CA
2.6±0.6	1.1	0.1
3.7±0.9	2.0	0.6
2.9±0.2	1.4	0.4
9.3±3.9	3.3	0.9

Table 3. Average annual home range size of three social groups of rhinoceros in Bardia.

Sex	Annual home range size			
	Mean	sd		
Single females (2)	22.1	3.7	$W_s = 8$	ns
Females with calf (6)	26.1	10.7		
Males (2)	41.8	5.1	$W_{s, m, f} = 19$	$p < 0.05$

Table 4. Seasonal home ranges of different social groups of Bardia animals in 1990-1993.

Sex	Hot			Monsoon			Winter			H	p
	n	\bar{X}	sd	n	\bar{X}	sd	n	\bar{X}	sd		
Females with small calves ^a	2	8.8	1.7	4	6.6	3.8	2	13.3	4.5	2.46	ns
Females with older calves ^b	4	14.4	3.2	3	18.6	11.2	3	20.9	9.4	2.17	ns
Single females	5	13.1	4.7	4	15.9	2.9	3	15.1	4.5	2.39	ns
Males	275: 29.0 _D			9.2 _{SD}			13.5 _{SD}				
	2	21.2	11.1	2	20.8	10.3	2	20.8	8.4	0.32	ns
	337: 13.3 _{SD}			30.4 _D			28.0 _D				

a = calves < 1 year, b = calves 1-5 years, D = dominant, SD = sub-dominant, H = Kruskal-Wallis test

Table 5. S

Seasons

Hot season

Monsoon s

Winter sea

All season

Females v

Wilcoxon

a animals in 1990-1993.

	Winter		H	p
	\bar{X}	sd		
2	13.3	4.5	2.46	ns
3	20.9	9.4	2.17	ns
3	15.1	4.5	2.39	ns
2	20.8	8.4	0.32	ns

ib-dominant, H = Kruskal-Wallis

Table 5. Seasonal changes in home ranges of females* in Bardia

Seasons	n	% change	t**	p
Hot season	5	+ 0.9	10	ns
Monsoon season	4	+38.1	4	ns
Winter season	2	-31.1	both decreasing	
All season	11	+ 8.6	41	ns

* Females with calves < 1 year excluded in comparison.
 ** Wilcoxon matched pair signed-rank test

Table 6. Mean percent overlap in seasonal home ranges (HR) and core areas (CA), mean distance between seasonal occupancy centers (km) and mean range lengths (km/48 hrs) for Bardia and Chitwan animals.

	Bardia				Chitwan			
	Females		Males		Females		Males*	
Seasonal overlap	HR	CA	HR	CA	HR	CA	HR	CA
Hot/Monsoon	74.8	43.8	51.3	21.6	65.6	27.6	75.3	9.8
Monsoon/Winter	54.9	21.6	41.2	20.4	67.1	29.4	79.1	44.7
Winter/hot	59.6	25.4	46.3	18.6	61.1	26.5	58.5	0
Distance between OC								
Hot/Monsoon	1.1±1.4		6.2±1.6		0.5±0.2		0.5	
Monsoon/winter	3.1±1.8		3.3±0.9		0.4±0.2		0.3	
Winter/hot	2.5±1.3		4.9±1.4		0.6±0.1		0.6	
Range length								
Hot	2.5±0.3		4.1±2.2		0.5±0.1		0.5	
Monsoon	2.9±1.4		2.3±1.3		0.4±0.2		0.4	
Winter	2.8±1.2		3.3±1.6		0.8±0.1		0.3	

* Only 1 male in Chitwan

Tabl
Nep

Habit
types

Sal f

Khair

River

Mixe

forest

Tall C

Wood

Phant

Bushy

Agricu

River

* differ

Areas (CA), mean distance
(18 hrs) for Bardia and

Chitwan			
Females		Males*	
HR	CA	HR	CA
65.6	27.6	75.3	9.8
67.1	29.4	79.1	44.7
61.1	26.5	58.5	0
0.5±0.2		0.5	
0.4±0.2		0.3	
0.6±0.1		0.6	
0.5±0.1		0.5	
0.4±0.2		0.4	
0.8±0.1		0.3	

Table 7. Utilization-availability data for habitat types in the Royal Bardia National Park, Nepal, 1990-93.

Habitat types	Total area km ²	Expected proportion of usage	Actual proportion of usage	Bonferroni intervals	Ivlev's Electivity Index
Sal forest	7.68	0.112	0.004	-0.001 ≤ p ≤ 0.009*	-0.93
Khair-Sissoo forest	13.54	0.197	0.411	0.376 ≤ p ≤ 0.446*	+0.35
Riverine forest	6.81	0.099	0.257	0.226 ≤ p ≤ 0.288*	+0.44
Mixed Hardwood forest	0.98	0.014	0.013	0.003 ≤ p ≤ 0.021	-0.07
Tall Grassland	8.19	0.119	0.218	0.189 ≤ p ≤ 0.247*	+0.29
Wooded Grassland	3.13	0.045	0.009	0.002 ≤ p ≤ 0.016*	-0.67
Phanta	0.95	0.014	0.0	0	0.0
Bushy Pasture	4.80	0.070	0.0	0	0.0
Agriculture	1.40	0.020	0.0	0	0.0
River and riverbeds	21.40	0.311	0.088	0.069 ≤ p ≤ 0.108*	-0.56

* difference at 0.05 level of significance.

Table 8. Habitat use by *Bardia rhinoceros* (% locations in each habitat) in hot, winter and monsoon seasons compared with the habitat composition of the study area.

Habitat types	% available	Hot		Monsoon		Winter	
		M	F	M	F	M	F
Sal forest	11.2	0	0.3 ^{ns}	0	1.2*	0	0
Khair-Sissoo forest	19.6	25.0 ^{ns}	35.1*	33.1 ^{ns}	30.5*	47.0*	50.7*
Riverine forest	9.9	32.7*	22.7*	16.7 ^{ns}	24.4*	41.2*	32.0*
Tall Grassland	11.9	19.2 ^{ns}	26.4*	36.3*	28.5*	11.8 ^{ns}	12.9 ^{ns}
Mixed Hardwood forest	1.4	1.9 ^{ns}	1.5 ^{ns}	2.8 ^{ns}	2.0 ^{ns}	0	0.4 ^{ns}
Wooded Grassland	4.5	0	1.4*	0	0.5*	0	0.9*
Phanta	1.4	0	0	0	0	0	0
Bushy Pasture	7.0	0	0	0	0	0	0
Cultivated land	2.0	0	0	0	0	0	0
River and riverbeds	31.1	21.2 ^{ns}	12.6*	11.1*	12.9*	0	3.1*

* difference at 0.05 level of significance.

at) a hot, winter study area.

Winter	
M	F
0	0
47.0*	50.7*
41.2*	32.0*
11.8 ^{ns}	12.9 ^{ns}
0	0.4 ^{ns}
0	0.9*
0	0
0	0
0	0
0	3.1*

Table 9. Composition (in percent) of habitats in the whole study area and in the seasonal core areas in Bardia in 1990-1993.

Habitat types	Composition of study area	Composition of core areas		
		Hot	Monsoon	Winter
Sal forest	11.2	0	0.1±0.2	0
Khair-Sissoo forest	19.6	30.3±2.9	28.6±4.0	40.4±5.4
Riverine forest	9.9	14.3±1.2	7.6±1.8	14.6±3.7
Tall Grassland	11.9	21.7±2.4	32.3±2.9	8.2±3.9
Mixed Hardwood forest	1.4	4.1±1.7	3.2±1.3	1.4±0.7
Wooded Grassland	4.5	4.1±0.9	4.0±1.5	9.0±3.6
Phanta	1.4	0	0.2±0.2	0
Bushy Pasture	7.0	0	0	0
Agriculture	2.0	0	0	0
River and riverbeds	31.1	25.5±1.7	24.0±1.3	26.4±0.8

sonal

er

4

7

9

6

8





Figure 1. Map of the western section of Royal Bardia National Park showing study area.



Inset map showing study area.




Figure 2. Seasonal home range and occupancy center of one adult female *Rhinoceros unicornis* in 1990/91.

Dotted line = monsoon (M), solid line = hot (H) and stipled line = winter (W). Floodplain consists of a mosaic of different habitat types (see text).




Figure 3. Range length of one adult female *Rhinoceros unicornis* during winter season, Royal Bardia National Park, 1990-1991.




Figure 6. Seasonal distribution of animal locations (in %) on three main habitat types in Bardia. * denotes significant preference by each sex (Bonferroni intervals) and non-matching letters denote a significant difference in habitat between sexes in each season.





Figure 5. Percent of locations in the individual home ranges compared with habitat composition of the home ranges. Filled circles are females, and open circles are males.



ange

as compared with habitat
and open circles are males.




Figure 4. Proportion of female locations in three habitat types in three seasons (χ^2 test: *, $p < 0.05$, ns = not significant)




Figure 7. Map of two study areas in Royal Bardia NP (middle section) and Royal Chitwan NP showing distribution of different habitat types.




Figure 7. Map of two study areas in Royal Bardia NP (middle section) and Royal Chitwan NP showing distribution of different habitat types.