Reproductive Disorders in White Rhinoceroses and the Value of Ultrasonographic Assessment

Hermes R., Hildebrandt T.B., Schwarzenberger F., Walzer C., Fritsch G. and F. Göritz Institute for Zoo Biology and Wildlife Research, Berlin, Germany

Based on endocrine monitoring 50% of the female white rhinoceros EEP population is considered to have no luteal activity. This lack of luteal activity represents one significant cause for the limited reproductive success of white rhinoceroses in captivity. Ultrasound evaluations of the reproductive tract combined with long-term endocrine monitoring were conducted in n=11 an-oestrous and 4 cycling females to elucidate the ovarian and uterine status. Five sonographic types of an-estrous (flat-lining) females were characterised: type I (n=3) small inactive ovaries, associated with uterine fluid accumulation in one older female; type II (n=2) persistent luteal structures; type III (n=1) animal with follicular cysts. type IV (n=4) young females with dynamic follicular development but lack of ovulation, type V (n=1) animal with massive para-ovarian tumour. In three regular cycling females cystic endometrial alteration was identified as predominant pathological finding. Due to the advanced status of alterations, a total of four females (≥27 years) were considered post-reproductive. Only females <11 years were free of pathological alterations. Reproductive tract ultrasound resulted in (1) the accurate evaluation of the breeding potential, (2) the sonographic characterisation of an-oestrous females and (3) the selection of candidates for natural or artificial breeding efforts. Increased use of ultrasonography in rhinoceros reproduction is useful tool to enhance breeding success in captivity.

Faecal Steroid Analysis in Free-Ranging Female White Rhinoceroses (Ceratotherium simum simum) on Otiiwa Game Ranch in Namibia

Mraz B¹, Modler F¹, Walzer C², Jago M³, Reinhardt P⁴, Schwarzenberger F¹,


¹Inst. Biochem. & LBI Vet. Med. Edocrinol., Vet med Univ, A-1210-Vienna, Austria

²Salzburg Zoo Hellbrunn, A-5081 Salzburg, Austria

³Otjiwarongo Vet. Clinic, Namibia

⁴Otjiwa Game Ranch, Namibia

Acyclicity is the major cause of low reproductive rates in captive white rhinoceroses: additional confounding factors are variable oestrous cycle lengths of 35 or 70 days and silent oestrus due to mate choice problems. The objectives of this study were to determine whether these conditions are present in wild populations. Study site was Otijwa game ranch, Namibia (100 km"; n=26 white rhinoceroses). Fresh dung samples were serially collected after tracking by game scouts. Samples were stored in methanol and dried prior to analysis with an established 20-oxo-pregnane assay. All females of breeding age (n=8) had calves; calving intervals of less than 2.5 years in the older animals (n=6) indicate that infertility conditions are not present in this wild population. Farm size and brush coverage made tracking difficult and average individual sample collection interval was 38.5 ± 5.0 d (range 1-100 d). This and the fact that females seem to become pregnant shortly after lactational anoestrus hindered statements on cyclicity. However, pregnancy diagnosis, even from single samples was 100% accurate after Day 120 (threshold value: 3000 ng/g faeces; significantly higher concentrations in >120d pregnant versus non-pregnant and <120d pregnant animals; p=0.001). In summary, pregnancy was accurately diagnosed by faecal steroid analysis in a wild white rhino population. Low reproductive rates, similar to those in the captive population were not present. For oestrous cycle diagnosis sample frequency during inter pregnancy periods needs to be increased. It can be concluded that faecal steroid analysis in wild white rhinoceros populations offers new perspectives for in situ and exsitu management of this species.



Harald M. Schwammer Thomas J. Foose Michael Fouraker Deborah Olson





A Research Update on Elephants and Rhinos

Proceedings of the International Elephant and Rhino Research Symposium, Vienna, June 7-11, 2001